

RESEARCH

Open Access



# Normal spirometry prediction equations for the Iranian population

Leyla Sahebi<sup>1</sup>, Besharat Rahimi<sup>2</sup>, Mamak Shariat<sup>1</sup>, Seyyed Hosein Mousavy<sup>3</sup> and Mohsen Hosseini<sup>4\*</sup>

## Abstract

**Background:** This study aimed to establish normative spirometric equations in a healthy population of Iranian children and adults, and compare these equations with those developed by the Caucasian Global Lung Initiative (GLI) for the first time.

**Methods:** Spirometric data from healthy Iranian aged 4–82 years sampled in 2019 were used to derive reference equations using the generalized additive model for location ( $\mu$ ), shape ( $\lambda$ ), and scale ( $\sigma$ ).

**Results:** A total of 418 females and 204 males were included in the study. Applying the GLI standards for the Iranian population resulted from the Z scores of FEV<sub>1</sub>, FVC, FEV<sub>1</sub>/FVC, and FEF<sub>25–75%</sub> was not different from zero. Based on the newly calculated LLN, eleven individuals showed significant values below the LLN for FEV<sub>1</sub>/FVC. In all age groups, this frequency was less than 5%, except for men over 70 years of age, which was 12.5%. There are significant differences between new data and GLI for Caucasian data.

**Conclusion:** It is recommended that the values and equations generated from this study should be used by physicians and technicians in their routine practice for the diagnosis and assessment of pulmonary disorders.

**Keywords:** Spirometry, Reference standards, Respiratory function tests

## Introduction

Experimental diagnosis of respiratory diseases, their intensity, and prognosis are principally dependent on spirometric results [1]. Accurate interpretation of spirometry requires standardized reference values that are predicted from its population race, as well as, age, and height [2–5].

In 2012, the Global Lung Function Initiative (GLI-2012) reported normative reference values, derived from over 160,000 data points in combined datasets from 33 countries. The GLI-2012 equations provided sex, age, height, and ethnic-specific reference equations as well as the lower limit of normal (LLN) values for spirometry

[5]. Although this approach included data from various countries, it did not include many populations.

Appropriateness of the GLI-2012 equations should be confirmed prior to their use for regions that are not currently covered by the reference equations [6]. In some studies has been confirmed suitability of GLI-2012 norms for their population, for example in the Australasian [7], Norwegian [8], German [9] and French [10] populations. But, the GLI-2012 norms seem inappropriate for use in the Swedish [11], Finnish [12], and Chinese [13] populations.

Our previous study found that Caucasian GLI equations were not suitable for the Iranian population, especially children under 10 years old [14]. The lack of predictive values specific to the Iranian population may lead to the misclassification of disease. Therefore, standardization of spirometry reference values is very necessary.

\*Correspondence: sorush140sr@gmail.com

<sup>4</sup> School of medicine, Tehran University of Medical Science, Tehran, Iran  
Full list of author information is available at the end of the article



Respiratory scales dedicated by spirometry to each person, do not follow a linear model, and the lung volume changes according to height and age with a skewed distribution. A practical approach that has been applied for spirometric data is Generalized Additive Models for Location, Scale, and Shape (GAMLSS). GAMLSS model is a non-parametric regression equation that best fits pulmonary and spirometric measures distribution. This model is the best existing one for the prediction of pulmonary values and the prediction equations offered by GLI-2012 have been confirmed and endorsed by many international respiratory societies. [15, 16].

Simulations show that when the confounders have a non-linear association with the outcome, compared to a parametric representation, GAMLSS modelling reduce the mean squared error for the adjusted exposure effect and avoid inflation of the type I error for testing the exposure effect [17].

This, was the first study in the Iranian population that aimed to predict the standard values of spirometry for Iranian reference population.

## Methods

### Design

This cross-sectional study was performed in Iran (Tehran) in 2019. This study was approved by the National Institute for Medical Research Development (NIMAD) (code: 978,931, 2019/05/28) and the Ethics Committee (code: IR.NIMAD.REC.1398.257). Conscious and written consent was obtained from all participants.

The study population was gathered from Tehran, and those who referred to local health centers- associated with Tehran Municipality- were included in the study. Overall 44 local health centers were selected by the randomized clustering method. The age range for inclusion in the study was 3–95 years. Informed consent was obtained from all participants / their legal guardians.

Healthy non-smokers between 3 and 95 years old, without a history of current airway or lung disease were included in the study. Exclusion criteria were as follows: not eligible for spirometry test and occurrence of respiratory disorders such as sputum cough, and rhinorrhea in the last 7 days.

Demographical and anthropometric variables such as sex, age, height, and weight were documented. Spirometric indices included FEV<sub>1</sub>, FVC, FEV<sub>1</sub>/FVC, and FEF<sub>25–75%</sub> (Forced expiratory flow averaged over the middle portion of FVC) were measured.

The validity, repeatability, and quality control were done according to the American Thoracic Society/European Respiratory Society (ATS/ERS) recommendations [18, 19], and described in more detail in an earlier paper (first phase of this study) [14].

In this study, 418 females and 204 males in different age groups (4–82 years old) were eligible to enter the study.

### Analysis

In the earlier study [14] we measured the lower limits of normal (LLN), Z-scores and percentiles for FEV<sub>1</sub>, FVC, FEV<sub>1</sub>/FVC, and FEF<sub>25–75%</sub> for each person. We determined agreement between the observed values in our population and the GLI reference values. According to the agreement by the GLI team, a mean Z-score outside the range of  $\pm 0.5$  was considered clinically significant [5, 11, 20, 21, 22]. The relationship between Z-scores and age, height, weight, and sex was examined using multiple linear regression models in the previous article [14].

In this study, GLI-2012 equations were implemented using the Generalized Additive Models for Location, Scale, and Shape (GAMLSS4.3-1) in software R (version 3.1.2; <http://www.r-project.org>), this method permitted the fitness of mean (M), coefficient of variance (S), and skewness (L) (Lambda–Mu–Sigma (LMS) of spirometry indices [5, 23].

The spirometry indices were modeled in males and females by age and height as explanatory variables using the Box-Cox-Cole–Green (BCG) distribution. The fit-test regression models were chosen by using Schwarz Bayesian Criterion (SBC), Akaike's Information Criterion (AIC) and assessing optimal degrees of freedom (df) for the cubic spline curve. The goodness of fit was also checked by normal Q–Q plots. Mean (M) indicates the predicted value as follows:  $M = \exp [a + b \times \ln(\text{height}_{\text{cm}}) + c \times \ln(\text{age}_{\text{year}}) + M\text{-spline}]$  (a, b, and c are coefficients, and M-spline is an age-specific contribution from the spline function. Values of L and S were also calculated based on regression output values of Sspline and Lspline. Finally we calculated LLN as follows:  $\text{LLN (5th percentage)} = \exp [\ln(M) + \ln(1 - 1.645 \times L \times S)/L]$ . Z-scores were calculated as (observed-predicted)/SD, where SD was calculated as (predicted-LLN)/1.645 [1, 23].

Agreement between Caucasian values and GLI-2012 Iranian prediction analyzed by Bland–Altman plots.

### Result

Six hundred and twenty-two Iranian participants (418 females and 204 males) aged 4–82 years were finally included in this study. The mean (range) age was 38.34 (4–82) years for men and 44.55 (4–80) years for women. The mean (SD) height for men and women were 1.72 (0.08) m and 1.58 (0.08) m over 21 years, respectively. Thirty-nine (19.2%) men and 131 (31.4%) women had a BMI  $\geq 30$  kg/m<sup>2</sup> (Table 1). Demographical and spirometry measurements of the reference population by gender are shown in Table 1 (Table 1).

**Table 1** Demographical and spirometry measurements of the reference population by gender

Variables	Females (414)	Males (206)
Age; mean (SD), y	45.54 (15.67)	37.78 (19.56)
Weight; mean (SD), Kg	68.61 (16.25)	74.88 (20.99)
Height; mean (SD), cm	156.18 (11.09)	168.22 (16.27)
BMI; mean (SD)	27.84 (5.74)	25.8 (5.23)
FEV <sub>1</sub> ; mean (SD), l	2.63 (0.65)	3.65 (1.04)
Min–Max	0.79–5.01	0.86–6.58
FVC; mean (SD), l	3.20 (0.78)	4.42 (1.24)
Min–Max	0.9–0.78	1.15–7.5
FEV <sub>1</sub> /FVC; mean (SD)	0.82 (0.70)	0.83 (0.067)
Min–Max	0.65–1.49	0.51–1.0
FEF <sub>25–75%</sub> ; mean (SD), l	3.88 (1.29)	1.51 (0.523)
Min–Max	0.49–8.13	0.55–2.22

BMI Body mass index, FEF<sub>25–75%</sub> Forced expiratory flow at 25 and 75% of the pulmonary volume, FEV<sub>1</sub> Forced expiratory volume in one second, FVC Forced expiratory vital capacity, Min–Max Minimum–Maximum, SD Standard deviation

The Caucasian GLI-2012 was applied to this sample in earlier study [14]. The mean Z-scores of FEV<sub>1</sub>, FVC and the FEV<sub>1</sub>/FVC for males and females in different age groups were higher than the Caucasian predicted values (range: 0.01 to 1.05) except for the FEV<sub>1</sub>/FVC in the age group under 21 years (range: –1.11 to –0.09).

The Z-scores of FEV<sub>1</sub>, FVC, FEV<sub>1</sub>/FVC, and FEF<sub>25–75%</sub> distribution based on Caucasian equation by sex and age in the Iranian healthy people is accessible in Table 2.

#### Iranian version of reference equations for spirometric values

We modeled GAMLSS regression equations for each spirometric parameter (FEV<sub>1</sub>, FVC, FEV<sub>1</sub>/FVC, and FEF<sub>25–75%</sub> obtained from the study population (Look up Tables and equations are available in the supplementary material). For new (Iranian) equations, none of the Z scores (FEV<sub>1</sub>, FVC, FEV<sub>1</sub>/FVC and FEF<sub>25–75%</sub>), are different from zero (by one-sample t-test analysis) in all age groups (P > 0.05). Distributions of Z-Scores based on Iranian (new) and Caucasian (GLI-2012 equations were shown in Table 2.

The obtained reference equations are used to estimate the Lower limit normal (LLN) of the spirometric parameters of FEV<sub>1</sub>, FVC, FEV<sub>1</sub>/FVC, and FEF<sub>25–75%</sub> for the Iranian population (Look up Tables are available in the supplementary material). Based on the newly calculated LLN, only five individuals showed significant values below the LLN for FEV<sub>1</sub>, FVC and eleven individuals for FEV<sub>1</sub>/FVC. This value was calculated in the Caucasian equation of 29 cases [14]. In all age groups, the frequency of Z-score for FEV<sub>1</sub>/FVC below the LLN was less than 5% except in men aged 70–84 years (12.5%) (Table 3). In the

Caucasian equations, the Z-score of FEV<sub>1</sub>/FVC was significantly higher among <21 years old (46.2% and 40.0% in males and females respectively). Frequency of FEV<sub>1</sub>/FVC < LLN by age and sex in Caucasian and Iranian equation is shown in Table 3.

Overall, residual Z-score for regression models was not beyond ±3 for our model (the standard range for residual is ±5). (Normal Q–Q plots (Additional file 1: Fig. S1a–h).

We found that age and height were the main predictors of the FEV<sub>1</sub> (males), FVC (males and females), and age for FEV<sub>1</sub>/FVC (not height) in both sex for final prediction models by nonlinear correlation analysis. The association between spirometric indices and anthropometric parameters is shown in Fig. 1a–l and Additional file 2: Table S1.

#### Agreement between Caucasian values and GLI-2012 Iranian prediction

The average differences (SDs) in FEV<sub>1</sub> (L), FVC and FEV<sub>1</sub>/FVC (%) predictions were –3.66(0.918), –3.7 (1.087), and –0.824 (0.034) for men, and –2.63 (0.49), –3.2(0.57), and –0.821 (0.003) for women, respectively. The Bland-Altman plots of FEV<sub>1</sub> (L), FVC and FEV<sub>1</sub>/FVC (%) are shown in Fig. 2a–f.

#### Discussion

This is the first study for the Iranian population that derived predictive equations and values using Lambda–Mu–Sigma (LMS) [18] by GAMLSS models. This model is preferable to the conventional multiple regression analysis which limits the model to several assumptions including normality of the residuals and constant variance [24]. On the other hand, LMS provides a variation in computing LLN through anthropometric data and prevents under-diagnosis of abnormalities in younger and taller individuals, and over-diagnosis of lung disorders in older and shorter people [25].

In this study, we have generated prediction equations for FEV<sub>1</sub>, FVC, FEV<sub>1</sub>/FVC, and FEF<sub>25–75%</sub> based on lung function data from 622 healthy Iranian populations. Genetic and environmental variables play a substantial role in the variability of lung function, so it is important to establish reference values appropriate to the ethnic and ecological characteristics of the local population [26, 27].

Our findings showed that GLI-2012 new equations adequately fitted FEV<sub>1</sub>, FVC, FEV<sub>1</sub>/FVC, and FEF<sub>25–75%</sub> data on the Iranian population for both genders.

In this population study of lung function, we assessed the agreement of lung function predictions between the GLI-2012 Caucasian values and GLI-2012 Iranian measures. The largest average difference was observed in FVC among men and the lowest difference was related to FEV<sub>1</sub>/FVC index in men and women.

**Table 2** Distributions of Z-Scores based on Iranian (new) and Caucasian (GLI-2012 equations (Assumption test: Compare means with zero)

Age group(Y)	Sex M	n	FEV1			FVC			FEV1/FVC			FEF <sub>25-75%</sub>							
			Mean ± SD		CI 95%	Mean ± SD		CI 95%	Mean ± SD		CI 95%	Mean ± SD		CI 95%					
			Caucasian	Iranian	Caucasian	Iranian	Caucasian	Iranian	Caucasian	Iranian	Caucasian	Iranian	Caucasian	Iranian					
<10	F	13	0.524 (0.621)	0.003 (0.197)	0.721 (0.627)	0.023 (0.214)	-1.11 (1.22)	-0.005 (0.14)	-0.111 (0.824)	-0.041 (1.013)	0.148, 0.899	-0.12, 0.122	0.341, 1.09	-0.107, 0.152	-1.84, -0.37	-0.087, 0.077	-0.609, 0.388	-0.65, 0.571	
			0.402 (0.904)	-0.33 (0.539)	0.618 (0.799)	-0.037 (0.534)	-1.21 (1.08)	-0.001 (0.12)	-0.589 (0.615)	-0.01 (0.7)	-0.589 (0.615)	-0.21, 0.825	-0.29, 0.22	0.244, 0.992	-0.287, 0.213	-1.72, -0.707	-0.061, 0.058	-0.878, 0.301	-0.338, 0.318
			0.269 (1.29)	0.008 (0.335)	0.396 (1.28)	0.011 (0.327)	-0.09 (1.01)	0.014 (0.13)	-0.089 (1.02)	0.044 (0.606)	-0.089 (1.02)	-0.155, 0.693	-0.102, 0.118	-0.026, 0.818	-0.119, 0.097	-0.421, 0.241	-0.06, 0.11	-0.425, 0.247	-0.155, 0.243
10-21	F	15	0.374 (1.086)	0.042 (0.327)	0.503 (0.685)	0.034 (0.198)	-0.323 (1.22)	0.022 (0.16)	-0.193 (1.46)	-0.005 (1.31)	-0.227, 0.975	-0.139, 0.223	0.124, 0.883	-0.0756, 0.143	-0.997, 0.35	-0.06, 0.11	-0.999, 0.614	-0.719, 0.729	
			0.162 (0.789)	-0.042 (0.208)	0.292 (0.714)	-0.03 (0.179)	0.101 (0.516)	0.009 (0.09)	-0.437 (0.946)	-0.0129 (0.598)	-0.193 (1.46)	-0.042 (0.208)	-0.137, 0.052	-0.032, 0.617	-0.112, 0.051	-0.135, 0.335	-0.029, 0.058	-0.867, 0.007	-0.401, 0.144
			0.145 (0.896)	0.022 (0.214)	0.423 (1.02)	0.018 (0.232)	0.014 (1.04)	0.007 (0.15)	-0.664 (1.1)	0.075 (0.685)	-0.242, 0.532	-0.071, 0.114	-0.019, 0.865	0.542 (1.22)	-0.083, 0.118	-0.436, 0.465	-0.05, 0.032	-1.097, 0.232	-0.221, 0.371
22-29	M	21	0.461 (1.27)	-0.005 (0.313)	0.542 (1.22)	-0.005 (0.297)	0.231 (0.688)	0.004 (0.11)	-0.106 (0.865)	0.029 (0.689)	0.069, 0.854	-0.101, 0.092	0.166, 0.917	-0.097, 0.086	0.019, 0.443	-0.05, 0.03	-0.444, 2.33	-0.183, 0.242	
			0.503 (1.24)	0.012 (0.315)	0.896 (1.29)	0.014 (0.313)	0.064 (0.665)	-0.006 (0.09)	-0.58 (0.865)	-0.008 (0.617)	0.069, 0.854	0.012 (0.315)	-0.056, 0.080	0.616, 1.18	-0.053, 0.082	-0.08, 0.209	-0.026, 0.013	-0.768 - 0.393	-0.142, 0.125
			0.234, 0.773	-0.056, 0.080	0.647 (0.935)	-0.027 (0.257)	0.420 (0.633)	0.005 (0.09)	-0.072 (0.769)	0.117 (0.591)	0.30, 0.996	-0.067, 0.132	0.261, 1.03	-0.079, 0.133	0.159, 0.681	-0.03, 0.039	-0.389, 2.46	-0.126, 0.361	
30-39	F	23	0.567 (1.16)	-0.039 (0.038)	0.906 (1.19)	-0.042 (0.357)	0.277 (0.774)	-0.006 (0.11)	-0.489 (1.08)	-0.057 (0.881)	0.567 (1.16)	-0.114, 0.036	0.667, 1.15	-0.113, 0.029	0.123, 0.431	-0.028, 0.017	-0.705 - 0.272	-0.233, 0.118	
			0.336, 0.797	-0.114, 0.036	0.618 (0.974)	0.062 (0.319)	0.417 (0.876)	0.028 (0.14)	-0.22 (0.888)	-0.127 (0.81)	0.336, 0.797	-0.041 (0.374)	0.618 (0.974)	0.618 (0.974)	0.417 (0.876)	0.034, 0.043	-0.558, 1.17	-0.432, 0.184	
			0.594 (1.09)	-0.102, 0.183	1.05 (1.48)	0.019 (0.396)	0.449 (0.784)	-0.005 (0.11)	-0.409 (0.905)	-0.0524 (0.954)	0.81, 1.01	-0.019 (0.395)	0.76, 1.33	-0.056, 0.096	0.298, 0.599	-0.011, 0.032	-0.558, 1.17	-0.131, 0.236	
40-49	M	25	0.615, 1.04	-0.02 (0.346)	0.301 (0.897)	-0.097 (0.293)	0.856 (0.838)	0.054 (0.13)	0.196 (1.078)	0.314 (1.344)	0.615, 1.04	-0.187, 0.147	-0.13, 0.733	0.452, 1.26	-0.08, 0.025	-0.323, 0.716	-0.333, 0.962		
			0.023, 1.01	-0.001 (0.581)	1.02 (1.18)	0.008 (0.569)	0.32 (0.757)	0.01 (0.11)	-0.533 (0.86)	-0.0425 (1.17)	0.023, 1.01	-0.001 (0.581)	0.723, 1.32	-0.135, 0.151	0.132, 0.513	-0.048, 0.010	-0.749 - 0.316	-0.338, 0.253	
			0.723 (1.15)	-0.147, 0.145	0.723, 1.32	-0.135, 0.151	0.132, 0.513	-0.096 (0.25)	0.04 (1.28)	-0.498 (1.73)	0.433, 1.01	-0.147, 0.145	0.476 (2.02)	-0.49, 0.32	-1.49, 0.269	-0.010, 0.118	-0.685, 0.676	-1.42, 0.426	
50-59	F	16	0.357 (2.23)	-0.68, 0.314	-0.599, 1.55	-0.49, 0.32	-1.49, 0.269	-0.010, 0.118	-0.685, 0.676	0.357 (2.23)	-0.68, 0.314	-0.599, 1.55	-0.49, 0.32	-1.49, 0.269	-0.010, 0.118	-0.685, 0.676	-1.42, 0.426		
			-0.829, 1.54	-0.68, 0.314	-0.599, 1.55	-0.49, 0.32	-1.49, 0.269	-0.010, 0.118	-0.685, 0.676	-0.829, 1.54	-0.68, 0.314	-0.599, 1.55	-0.49, 0.32	-1.49, 0.269	-0.010, 0.118	-0.685, 0.676	-1.42, 0.426		
			0.356 (0.822)	-0.12 (0.495)	0.06 (0.662)	-0.152 (0.385)	0.073 (0.983)	-0.016 (0.16)	0.085 (1.08)	0.009 (1.93)	0.356 (0.822)	-0.12 (0.495)	0.06 (0.662)	-0.152 (0.385)	0.073 (0.983)	-0.016 (0.16)	0.085 (1.08)	0.009 (1.93)	

**Table 2** (continued)

Age group(Y)	Sex M	n	FEV1		FVC		FEV1/FVC		FEF <sub>25-75%</sub>	
			Mean ± SD	CI 95%	Mean ± SD	CI 95%	Mean ± SD	CI 95%	Mean ± SD	CI 95%
	Caucasian	Iranian			Caucasian	Iranian	Caucasian	Iranian	Caucasian	Iranian
			- 0.331, 1.043	- 0.532, 0.295	- 0.494, 0.614	- 0.475, 0.169	- 0.748, 0.896	- 0.12, 0.149	- 0.819, 989	- 1.61, 0.624


F = Female, M = male

FEV<sub>1</sub> Forced expiratory volume, FEF<sub>25-75%</sub> Forced expiratory flow at 25 and 75% of the pulmonary volume, FEV<sub>1</sub> Forced expiratory volume in one second, FVC Forced expiratory vital capacity, SD Standard deviation

**Table 3** Frequency of FEV<sub>1</sub>/FVC < LLN by age and sex in Caucasian and Iranian equation

Age groups	Total (622)			Males (204)			Females (418)		
	n	Caucasian	Iranian	n	Caucasian	Iranian	n	Caucasian	Iranian
< 10	33	14 (42.4)	0	13	6 (46.2)	0	20	8 (40.0)	0
10–21	53	4 (7.5)	2 (3.78)	38	2 (5.3%)	1 (2.63)	15	2 (13.3)	1 (6.7)
22–29	44	2 (4.5)	0	21	0	0	23	2 (8.7)	0
30–39	127	1 (0.8)	1 (0.8)	43	0	0	84	1 (1.2)	1 (1.2)
40–49	124	1 (0.8)	1 (0.8)	25	0	0	99	1 (1.0)	1 (1.0)
50–59	135	3 (2.2)	2 (1.15)	29	1 (3.4)	1 (3.4)	106	2 (1.9)	1 (0.94)
60–69	82	0	3 (3.66)	19	0	0	63	0	3 (4.8)
70–84	24	4 (16.7)	2 (8.3)	16	4 (25.0)	2 (12.5)	8	0	0

LLN Lower limit of normal



**Fig. 1** Nonlinear correlation analysis between FEV<sub>1</sub>, FVC and FEV<sub>1</sub>/FVC, and anthropometric parameters


In a study conducted on Jordanian people over 18 years old, based on Bland and Altman results, there were significant differences between the new equation and GLI for Caucasians equations too [24, 28].

In our study, age and height was the main predictors of the FEV<sub>1</sub> (males), FVC (males and females), and age for FEV<sub>1</sub>/FVC (not height) for both sexes.

In different similar studies on various ethnicities, anthropometric predictors have been measured on spirometric indices in both sexes. In a study conducted in India, it was found that age and height were the main predictors of the FEV<sub>1</sub> and FVC spirometry parameters

in both sexes, for FEV<sub>1</sub>/FVC, only age was a significant predictor of outcome [29] but not height. This result was consistent with the findings of our study. Chang’s and colleagues reported, the height and weight, but not age, were important predictors in the final prediction models for FVC and FEV<sub>1</sub> in Taiwanese children [30].

In our study, the frequency Z-score of FEV<sub>1</sub>/FVC below LLN was less than 5% in all age groups, except for the group of men over 70 years old (12.5%). This finding was consistent with the results of a study conducted in India [29]. But this amount was estimated at 10% in Mozambique’s reference population (Southeast Africa) [31],



also the LLNs of FEV<sub>1</sub>/FVC were less than 0.70 in men above 56 years of age and women above 60 years of age in Chinese aged 4–80 years [1]. Concerning the high prevalence in men over 70 years of age, this may be due to the low sample size in this age group (16 people). However, the initial interview to enter the study was accompanied strictly, but the possibility of bias could not be prevented absolutely. For example, some elderly men may have had the experience of smoking in the past but have forgotten or for some reason declare that they have not had this experience.

This study has several limitations. First, the sample size of this study is not very large. However, we would claim that the sample size of men and women is large enough to have enough power for validating spirometry reference values (at least 150 subjects for each gender) [32].

**Conclusion**

GLI-2012 Iranian equations fitted FEV<sub>1</sub>, FVC, FEV<sub>1</sub>/FVC, and FEF<sub>25–75%</sub> data of Iranian population for both gender. There were significant differences between measures by GLI for Caucasians and Iranian (new) equations. It is recommended that the values and equations generated from this study should be used by physicians and experts in practice for detecting the disease condition and its severity in Iranian populations.

**Supplementary Information**

The online version contains supplementary material available at <https://doi.org/10.1186/s12890-022-02273-8>.

**Additional file 1: Fig. S1.** Normal Q–Q plots for FEV<sub>1</sub>, FVC, FEV<sub>1</sub>/FVC and FEF<sub>25–75%</sub> by gender.

**Additional file 2: Table S1.** The association between spirometric indices and anthropometric parameters.

**Acknowledgements**

The authors wish to thank the staff of health houses affiliated with Tehran Municipality and all participants for their collaboration with the research team.

**Author contributions**

LS, BR: Writing proposal. BR, LS, SHM: data gathering. SHM: Administration Manager. BR: Clinical manager. LS and MH: Statistical analysis. LS, MSH: Writing draft. BR: Editing manuscript. All authors read and approved the final manuscript.

**Funding**

Noting.

**Availability of data and materials**

The datasets generated and/or analyzed during the current study are available from Dr. Leyla Sahebi (first author) on reasonable request.

**Declarations**

**Ethical approval and consent to participate**

This article was the result of a research project approved by the National Institute for Medical Research Development (NIMAD) (code: 978931, 2019/05/28) and the Ethics Committee (code: IR.NIMAD.REC.1398.257). All methods were carried out in accordance with relevant guidelines and regulations. Written

and informed consent was obtained from all participants for pulmonary function test using spirometry.

#### Consent for publication

No personal information of the participants in the article was reported.

#### Competing interests

The authors declare that they have no competing interests.

#### Author details

<sup>1</sup>Maternal, Fetal and Neonatal Research Center, Family Health Research Institute, Tehran University of Medical Sciences, Tehran, Iran. <sup>2</sup>Advanced Thoracic Research Center, Tehran University of Medical Sciences, Tehran, Iran. <sup>3</sup>Municipality of Tehran, Tehran, Iran. <sup>4</sup>School of medicine, Tehran University of Medical Science, Tehran, Iran.

Received: 15 August 2022 Accepted: 2 December 2022

Published online: 12 December 2022

#### References

- Jian W, Gao Y, Hao C, et al. Reference values for spirometry in chinese aged 4–80 years. *J Thorac Dis*. 2017;9(11):4538–49. <https://doi.org/10.21037/jtd.2017.10.110>.
- Braun L, Wolfgang M, Dickersin K. Defining race/ethnicity and explaining difference in research studies on lung function. *Eur Respir J*. 2013;41(6):1362–70. <https://doi.org/10.1183/09031936.00091612>.
- Strippoli MP, Kuehni CE, Dogaru CM, et al. Etiology of ethnic differences in childhood spirometry. *Pediatrics*. 2013;131(6):e1842–9. <https://doi.org/10.1542/peds.2012-3003>.
- Yang TS, Peat J, Keena V, Donnelly P, Unger W, Woolcock A. A review of the racial differences in the lung function of normal Caucasian, Chinese and Indian subjects. *Eur Respir J*. 1991;4(7):872–80.
- Quanjer PH, Stanojevic S, Cole TJ, et al. Multi-ethnic reference values for spirometry for the 3–95-yr age range: the global lung function 2012 equations. *Eur Respir J*. 2012;40(6):1324–43. <https://doi.org/10.1183/09031936.00080312>.
- Blake TL, Chang AB, Chatfield MD, Marchant JM, McElrea MS. Global lung function Initiative-2012 'other/mixed' spirometry reference equation provides the best overall fit for Australian Aboriginal and/or Torres Strait Islander children and young adults. *Respirology (Carlton Vic)*. 2020;25(3):281–8. <https://doi.org/10.1111/resp.13649>.
- HALL GL, THOMPSON BR, et al. The global lung Initiative 2012 reference values reflect contemporary Australasian spirometry. *Respirology (Carlton Vic)*. 2012;17(7):1150–1. <https://doi.org/10.1111/j.1440-1843.2012.02232.x>.
- Langhammer A, Johannessen A, Holmen TL, et al. Global lung function Initiative 2012 reference equations for spirometry in the norwegian population. *Eur Respir J*. 2016;48(6):1602–11. <https://doi.org/10.1183/13993003.00443-2016>.
- Hüls A, Krämer U, Stolz S, et al. Applicability of the global lung Initiative 2012 reference values for spirometry for longitudinal data of elderly women. *PLoS One*. 2016;11(6):e0157569. <https://doi.org/10.1371/journal.pone.0157569>.
- Hulo S, de Broucker V, Giovannelli J, et al. Global lung function Initiative reference equations better describe a middle-aged, healthy french population than the European community for steel and coal values. *Eur Respir J*. 2016;48(6):1779–81. <https://doi.org/10.1183/13993003.00606-2016>.
- Backman H, Lindberg A, Sovijärvi A, Larsson K, Lundbäck B, Rönmark E. Evaluation of the global lung function initiative 2012 reference values for spirometry in a swedish population sample. *BMC Pulm Med*. 2015;25:26. <https://doi.org/10.1186/s12890-015-0022-2>.
- Kainu A, Timonen KL, Toikka J, et al. Reference values of spirometry for finnish adults. *Clin Physiol Funct Imaging*. 2016;36(5):346–58. <https://doi.org/10.1111/cpf.12237>.
- Zhang J, Hu X, Shan G. Spirometry reference values for population aged 7–80 years in China. *Respirology (Carlton Vic)*. 2017;22(8):1630–6. <https://doi.org/10.1111/resp.13118>.
- Sahebi L, Rahimi B, Shariat M, Mousavy SH. Evaluation of the Global Lung Function Initiative 2012 reference values for spirometry in an Iranian population. *Sci Rep*. 2022;12(1):12784. <https://doi.org/10.1038/s41598-022-17306-9>.
- Cole TJ, Stanojevic S, Stocks J, Coates AL, Hankinson JL, Wade AM. Age- and size-related reference ranges: a case study of spirometry through childhood and adulthood. *Stat Med*. 2009;28(5):880–98. <https://doi.org/10.1002/sim.3504>.
- Rigby RA, Stasinopoulos DM. Smooth centile curves for skew and kurtotic data modelled using the Box-Cox power exponential distribution. *Stat Med*. 2004;15(19):3053–76. <https://doi.org/10.1002/sim.1861>.
- Benedetti A, Abrahamowicz M. Using generalized additive models to reduce residual confounding. *Stat Med*. 2004;30(24):3781–801. <https://doi.org/10.1002/sim.2073>.
- Pellegrino R, Viegi G, Brusasco V, et al. Interpretative strategies for lung function tests. *Eur Respir J*. 2005;26(5):948–68. <https://doi.org/10.1183/09031936.05.00035205>.
- Hankinson JL, Eschenbacher B, Townsend M, Stocks J, Quanjer PH. Use of forced vital capacity and forced expiratory volume in 1 second quality criteria for determining a valid test. *Eur Respir J*. 2015;45(5):1283–92. <https://doi.org/10.1183/09031936.00116814>.
- Abdullah N, Borhanuddin B, Shah SA, Hassan T, Jamal R. Global Lung Initiative 2012 spirometry reference values in a large Asian cohort of Malay, Chinese and Indian ancestry. *Respirology (Carlton, Vic)*. 2018;23(12):1173–9. <https://doi.org/10.1111/resp.13330>.
- Ben Saad H, El Attar MN, Hadj Mabrouk K, et al. The recent multi-ethnic Global Lung Initiative 2012 (GLI2012) reference values don't reflect contemporary adult's north African spirometry. *Respirology (Carlton, Vic)*. 2013;107(12):2000–8. <https://doi.org/10.1016/j.rmed.2013.10.015>.
- Arigliani M, Canciani MC, Mottini G, et al. Evaluation of the Global Lung Initiative 2012 reference values for spirometry in African children. *Am J Respir Crit Care Med*. 2017;195(2):229–36. <https://doi.org/10.1164/rccm.201604-0693OC>.
- Quanjer PHSS, Stocks J, Cole TJ, Stocks J. GAMLSS in action, 2012.
- Al-Qerem W, Hammad A, Gassar E, Al-Qirir M, Ling J. Spirometry reference equations for an adult Middle Eastern population. *Expert Rev Respir Med*. 2019. <https://doi.org/10.1080/17476348.2019.1601560>.
- Culver BH. How should the lower limit of the normal range be defined? *Respir Care*. 2012;57(1):136–45. <https://doi.org/10.4187/respcare.01427>. discussion 143–5.
- Schwartz J, Katz SA, Fegley RW, Tockman MS. Sex and race differences in the development of lung function. *Am Rev Respir Dis*. 1988;138(6):1415–21. <https://doi.org/10.1164/ajrccm/138.6.1415>.
- Quanjer PH, Stanojevic S. Do the global lung function Initiative 2012 equations fit my population? *Eur Respir J*. 2016;48(6):1782–5. <https://doi.org/10.1183/13993003.01757-2016>.
- Al-Qerem W, Allassi A, Jarab A, Ling J. The applicability of the Global Lung Initiative equations and other regional equations on a sample of healthy Middle Eastern adolescents. *Clin Respir J*. 2021. <https://doi.org/10.1111/crj.13329>.
- Agarwal D, Parker RA, Pinnock H, et al. Normal spirometry predictive values for the western indian adult population. *Eur Respir J*. 2020. <https://doi.org/10.1183/13993003.02129-2019>.
- Chang SM, Tsai HJ, Tzeng JY, et al. Reference equations for spirometry in healthy asian children aged 5 to 18 years in Taiwan. *World Allergy Org J*. 2019;12(11):100074. <https://doi.org/10.1016/j.waojou.2019.100074>.
- Ivanova O, Khosa C, Bakuli A, et al. Lung function testing and prediction equations in adult population from Maputo, Mozambique. *Int J Environ Res Public Health*. 2020. <https://doi.org/10.3390/ijerph17124535>.
- Quanjer PH, Stocks J, Cole TJ, Hall GL, Stanojevic S. Influence of secular trends and sample size on reference equations for lung function tests. *Eur Respir J*. 2011;37(3):658–64. <https://doi.org/10.1183/09031936.00110010>.

#### Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.